ELSEVIER

Journal of Immunological Methods 219 (1998) 197–198

JOURNAL OF IMMUNOLOGICAL METHODS

Contents of volume 219 (1998)

Numbers 1 and 2 – 1 October 1998

Immunological standardization

Cytokine immunoassays: recommendations for standardisation, calibration and validation
M. Wadhwa and R. Thorpe (UK) 1

Research papers

Measurement and biological correlates of antibody bioactivity during antibody immunotherapies
D.S. Hagg and R.P. Junghans (USA) 7

A sandwich type Acridinium-Linked ImmunoSorbent Assay (ALISA) detects soluble ErbB1 (sErbB1) in normal human sera
A.T. Baron, J.M. Lafky, D.C. Connolly, J. Peoples, D.J. O'Kane, V.J. Suman, C.H. Boardman, K.C. Podratz and N.J. Maihle (USA) 23

A sensitive fluorometric enzyme-linked immunosorbent assay that measures vascular endothelial growth factor165 in human plasma
C.R. Rodriguez, D.T. Fei, B. Keyt and D.L. Baly (USA) 45

Green- and red-fluorescent nanospheres for the detection of cell surface receptors by flow cytometry
M.K. Bhalgat, R.P. Haugland, J.S. Pollack, S. Swan and R.P. Haugland (USA) 57

Optimisation of the conditions for generating human DC initiated antigen specific T lymphocyte lines in vitro
S.I. Mannering, J.L. McKenzie and D.N.J. Hart (New Zealand) 69

A new in vitro model for studying human T cell differentiation: TH1/TH2 induction following activation by superantigens
S. Gehring, M. Schlaak and J. Van der Bosch (Germany) 85

Variability between methods to determine ANA, anti-dsDNA and anti-ENA autoantibodies: a collaborative study with the biomedical industry
N. Bizzaro, R. Tozzoli, E. Tonutti, A. Piazza, F. Manoni, A. Ghirardello, D. Bassetti, D. Villalta, M. Pradella and P. Rizzotti (Italy) 99

A simplified procedure for the isolation of immunoglobulins from human serum using a novel type of thiophilic gel at low salt concentration
G.H. Scholz, S. Vieweg, S. Leistner, J. Seissler, W.A. Scherbaum and K. Huse (Germany) 109

Highly efficient recovery of functional single-chain Fv fragments from inclusion bodies overexpressed in *Escherichia coli* by controlled introduction of oxidizing reagent—application to a human single-chain Fv fragment
K. Tsumoto, K. Shinoki, H. Kondo, M. Uchikawa, T. Juji and I. Kumagai (Japan) 119

Detection of HLA-B27 alleles by group-specific amplification and time-resolved fluorometry
L. Välimaa, M. Sjöroos, M. Luhtala, P. Toivanen, T. Lövgren and J. Ilonen (Finland) 131

New ligands for HLA DRB1*0301 by random selection of favourable amino acids ranked by competition studies with undecapeptide amide sublibraries
C. Jung, M. Kalbus, B. Fleckenstein, A. Melms, G. Jung and K.-H. Wiesmüller (Germany) 139

Monoclonal antibodies raised to paraffin wax embedded archival tissue; feasibility study of their potential to detect novel antigenic markers
E. Moran, A. Larkin, I. Cleary, C. Barnes, S.M. Kennedy, P. Kelehan and M. Clynes (Ireland) 151

Variability in the sensitivity of nine enzyme-linked immunosorbant assays (ELISAs) in the measurement of human interleukin 6
T. Krakauer (USA) 161

RT-PCR quantitation of cytokine responses in vivo from specimens containing small numbers of cells during bioimmunotherapy
M.A. Nash, R. Lenzi, C.D. Platsoucas and R.S. Freedman (USA) 169

A flow cytometric assay for D8/17 B cell marker in patients with Tourette's syndrome and obsessive compulsive disorder
F. Chapman, K. Visvanathan, R. Carreño-Manjarrez and J.B. Zabriskie (USA) 181

Short communication

A simple flow cytometry assay using dihydrorhodamine for the measurement of the neutrophil respiratory burst in whole blood: comparison with the quantitative nitrobluetetrazolium test
M.P. Richardson, M.J. Ayliffe, M. Helbert and E.G. Davies (UK) 187

Announcement 195

Contents of volume 219 (1998) 197

Notes to Authors 199

Protocol: Notes to Authors 201